

**Reported Communicable Diseases, North Carolina, January-December 2010 (by date of report)\***

Disease	Year-to-Date (Fourth Quarter)			4thQuarter 2010	Comments / Note
	2010	2009	Mean (2005-2009)		
Brucellosis	1	1	3	0	
Campylobacter	851	587	656	267	
Chlamydia, laboratory reports	45,190	43,734	35,406	11,100	
Cholera	1	0	0	1	
Cretzfeldt-Jacob Disease	13	13	4	7	
Cryptosporidiosis	94	160	115	21	
Cyclosporiasis	1	2	2	1	
Dengue	6	0	7	1	
E. coli Shiga Toxin-producing	97	112	122	51	
Ehrlichiosis, Granulocytic	36	3	1	11	Note 1
Ehrlichiosis, Monocytic	148	73	47	20	Note 1
Gonorrhea	15,125	14,811	15,774	3,785	
Haemophilus Influenzae	123	105	74	19	
Hepatitis A	48	41	69	7	
Hepatitis B	115	104	138	32	
Hepatitis B Carrier	693	898	865	267	
Hepatitis B Perinatal	1	1	2	1	
Hepatitis C, Acute	39	24	24	7	
HIV/AIDS	1,620	1,791	2,009	327	Note 2
Hemol.Urem.Syn/TTP	6	4	6	2	
Influenza, Adult death	25	75	N/A	0	
Influenza, Pediatric death	0	9	2	0	
Lacrosse (California)	23	169	34	12	
Legionellosis	63	65	47	13	
Leprosy (Hansen's Disease)	1	0	0	0	
Leptospirosis	1	0	1	0	
Listeriosis	22	27	28	6	
Lyme Disease	417	252	58	44	Note 1
Malaria	46	33	32	11	
Meninccocal Invasive Disease	14	32	27	2	
Meningitis, Pneumococcal	31	29	35	8	
Q Fever	1	1	3	0	
Rabies in animals	397	473	476	81	
RMSF	505	325	587	115	Note 1
Salmonellosis	2,346	1,806	1,739	1,246	
Shigellosis	254	358	228	116	
Strep A Invasive Disease	146	107	138	23	
Syphilis, Total	728	937	621	168	Note 3
Toxic Shock Synd.,Strep	8	4	7	0	
Toxic Shock Synd, non-Strep	2	0	4	1	
Tuberculosis	296	250	326	103	
Tularemia	3	1	1	0	
Typhoid, Acute	9	5	6	4	
Vaccinia	1	1	N/A	0	
V Vulnificus	7	3	4	0	
Vibrio, Other	17	12	14	7	
VISA/VRSA (Staph aureus)	3	2	1	1	
Whooping Cough	153	216	250	17	

\* Preliminary data, as of 01/01/2011. Quarters defined as 13-week periods. Diseases reported in 2010 define those listed in this table. Notes: 1. Suspect, Probable and Confirmed cases reported for 2009 and 2010. The five-year average for these events is based on confirmed and probable classifications only, because suspect classifications were not reportable prior to 2008; 2. Earliest report with HIV infection or AIDS diagnosis; 3. Includes primary, secondary and early latent syphilis.